

A NEW VARIETY OF PENSTEMON

Ralph W. Bennett

In *Phytologia*, Volume 9, Number 2, pp. 57--58, 1963, certain nomenclatural changes and new varieties were published. The following new variety is published herewith:

Penstemon Mitch.

P. hirsutus Willdenow

var. minimus Bennett, var. nov.

A var. hirsuti differt: caulibus non altior quam 10 cm.

- - - - -

NOTES ON NEW AND NOTEWORTHY PLANTS. XLII

Harold N. Moldenke

ALOYSIA BARBATA var. ACAPULCENSIS Moldenke, var. nov.

Haec varietas a forma typica speciei foliis ellipticis usque 11.5 cm. longis 4.5 cm. latis subtus minute puberulis resinoso-punctatisque recedit.

This variety differs from the typical form of the species in having its leaf-blades elliptic, to 11.5 cm. long, 4.5 cm. wide, and minutely puberulent and resinous-punctate beneath.

The type of the variety was collected by Ladislao Paray (no. 2666) at Puerto Matqués, near Acapulco, Guerrero, Mexico, on December 23, 1957, and is deposited in the herbarium of the Escuela Nacional de Ciencias Biologicas, Instituto Politecnico Nacional, in Mexico City.

CLERODENDRUM INTERMEDIUM f. ALBIFLORUM Moldenke, f. nov.

Haec forma a forma typica speciei corollis albis recedit.

This form differs from the typical form of the species in having white corollas.

The type of the form was collected by Francesco Fuerrero at Arayat, Panpanga Province, Luzon, Philippine Islands, in 1927, and is deposited in the herbarium of the University of California at Berkeley.

CLERODENDRUM LINDLEYI f. ALBIFLORUM Moldenke, f. nov.

Haec forma a forma typica speciei corollis albis recedit.

This form differs from the typical form of the species in having white corollas.

The type of the form was collected by Tsiang Ying (no. 1433) in open scrub on the way to Sie-kun, Lokchong Hsien, in the North